

Result summary

# Microdioriet; steenslag en brekerzand 2022 incl. belading vrachtauto

De Hoop Bouwgrondstoffen B.V.

Calculation number:	EPD-NIBE-20221120-32063
Generation on:	31-01-2023
Issue date:	31-01-2023
Valid until:	31-01-2028
Status:	verified

R<THiNK



## 1 General information

### 1.1 PRODUCT

Microdioriet; steenslag en brekerzand 2022 incl. belading vrachtauto



### 1.2 VALIDITY

Issue date 31-01-2023

Valid until: 31-01-2028

Msc. P.F. Stadhouders, EcoReview V.O.F.

### 1.3 OWNER OF THE DECLARATION



**Manufacturer:** De Hoop Bouwgrondstoffen B.V.

**Address:** Duitslandweg 2, 4538 BK Terneuzen

**E-mail:** info@dehoop-bouwgrondstoffen.nl

**Website:** www.dehoop-bouwgrondstoffen.nl

**Production location:** Carrières Unies de Porphyre

**Address production location:** Chemin de Mons a Gand, 7860 Lessines

### 1.4 VERIFICATION OF THE DECLARATION

CEN standard EN 15804 serves as the core PCR. In compliance with ISO 14040:2006 and 14044:2006.

Independent verification of the declaration according to EN ISO 14025:2011-10.

Internal  External

### 1.5 THIS DECLARATION IS BASED ON THE PRODUCT CATEGORY RULES

NMD Determination method Environmental performance Construction works v1.1 March 2022

### 1.6 FUNCTIONAL / DECLARED UNIT

Declared unit: ton (ton)

### 1.7 CONVERSION FACTORS

Description	Value	Unit
Declared unit	1	ton
Weight per declared unit	1000.000	kg
Conversion factor to 1 kg	0.001000	ton

### 1.8 SCOPE OF DECLARATION AND SYSTEM BOUNDARIES

This is a Cradle to gate LCA. The life cycle stages included are as shown below:

(X = module included, ND = module not declared)

A1	A2	A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
X	X	X	ND													

## 1 General information

### 1.9 COMPARABILITY

In principle, a comparison or assessment of the environmental impacts of different products is only possible if they have been prepared in accordance with EN 15804. For the evaluation of the comparability, the following aspects have to be considered in particular:

PCR used, functional or declared unit, geographical reference, the definition of the system boundary, declared modules, data selection (primary or secondary data, background database, data quality), scenarios used for use and disposal phases, and the life cycle inventory (data collection, calculation methods, allocations, validity period). PCRs and general program instructions of different EPDs programs may differ. Comparability needs to be evaluated. For further guidance, see EN 15804+A2 (5.3 Comparability of EPD for construction products) and ISO 14025 (6.7.2 Requirements for comparability).

## 2 Product

### 2.1 PRODUCT DESCRIPTION

Deze LCA geeft de milieueffecten weer van de productie van Microdioriet steenslag en brekerzand (in alle mogelijke sorteringen in de maten vanaf 0 tot en met 31,5 mm) Beschouwd zijn de processen in de groeve, intern transport en het beladen van vrachtwagens

Adres van de groeve:

Carrières Unies de Porphyre Chaussée Maieur Habilis 177, 1430 Bierghes, België



### 2.2 DESCRIPTION PRODUCTION PROCESS

Microdioriet wordt bij Carrières Unies de Porphyre te Bierghes (B) geproduceerd. De steen wordt met behulp van explosieven uit het massief gesprongen. De steen wordt daarna gebroken en gezeefd.

## 2 Product

Leeswijzer:

Alle processen in A1, A2 en A3 zijn opgenomen in A3

Alle processen in de groeve, brekerij , intern transport en beladen transportmiddel voor levering aan afnemers zijn beschouwd in A1+A2+A3.

## 3 Results

### 3.1 ENVIRONMENTAL IMPACT INDICATORS PER TON

#### CORE ENVIRONMENTAL IMPACT INDICATORS EN15804+A2

Abbreviation	Unit	A1	A2	A3	Total
AP	mol H+ eqv.	0.00E+0	0.00E+0	3.77E-2	3.77E-2
GWP-total	kg CO <sub>2</sub> eqv.	0.00E+0	0.00E+0	4.54E+0	4.54E+0
GWP-b	kg CO <sub>2</sub> eqv.	0.00E+0	0.00E+0	9.97E-3	9.97E-3
GWP-f	kg CO <sub>2</sub> eqv.	0.00E+0	0.00E+0	4.52E+0	4.52E+0
GWP-luluc	kg CO <sub>2</sub> eqv.	0.00E+0	0.00E+0	4.65E-3	4.65E-3
EP-m	kg N eqv.	0.00E+0	0.00E+0	1.39E-2	1.39E-2
EP-fw	kg P eqv.	0.00E+0	0.00E+0	7.55E-5	7.55E-5
EP-T	mol N eqv.	0.00E+0	0.00E+0	1.62E-1	1.62E-1
ODP	kg CFC 11 eqv.	0.00E+0	0.00E+0	7.72E-7	7.72E-7
POCP	kg NMVOC eqv.	0.00E+0	0.00E+0	4.11E-2	4.11E-2
ADP-f	MJ	0.00E+0	0.00E+0	8.75E+1	8.75E+1
ADP-mm	kg Sb-eqv.	0.00E+0	0.00E+0	2.28E-4	2.28E-4
WDP	m <sup>3</sup> world eqv.	0.00E+0	0.00E+0	8.82E-1	8.82E-1

**AP**=Acidification (AP) | **GWP-total**=Global warming potential (GWP-total) | **GWP-b**=Global warming potential - Biogenic (GWP-b) | **GWP-f**=Global warming potential - Fossil (GWP-f) | **GWP-luluc**=Global warming potential - Land use and land use change (GWP-luluc) | **EP-m**=Eutrophication marine (EP-m) | **EP-fw**=Eutrophication, freshwater (EP-fw) | **EP-T**=Eutrophication, terrestrial (EP-T) | **ODP**=Ozone depletion (ODP) | **POCP**=Photochemical ozone formation - human health (POCP) | **ADP-f**=Resource use, fossils (ADP-f) | **ADP-mm**=Resource use, minerals and metals (ADP-mm) | **WDP**=Water use (WDP)

#### ADDITIONAL ENVIRONMENTAL IMPACT INDICATORS EN15084+A2

Abbreviation	Unit	A1	A2	A3	Total
ETP-fw	CTUe	0.00E+0	0.00E+0	5.47E+3	5.47E+3
PM	disease incidence	0.00E+0	0.00E+0	9.17E-7	9.17E-7
HTP-c	CTUh	0.00E+0	0.00E+0	3.47E-9	3.47E-9
HTP-nc	CTUh	0.00E+0	0.00E+0	8.35E-8	8.35E-8
IR	kBq U235 eqv.	0.00E+0	0.00E+0	7.05E-1	7.05E-1

**ETP-fw**=Ecotoxicity, freshwater (ETP-fw) | **PM**=Particulate Matter (PM) | **HTP-c**=Human toxicity, cancer (HTP-c) | **HTP-nc**=Human toxicity, non-cancer (HTP-nc) | **IR**=Ionising radiation, human health (IR) | **SQP**=Land use (SQP)

### 3 Results

Abbreviation	Unit	A1	A2	A3	Total
SQP	Pt	0.00E+0	0.00E+0	7.75E+1	7.75E+1

**ETP-fw**=Ecotoxicity, freshwater (ETP-fw) | **PM**=Particulate Matter (PM) | **HTP-c**=Human toxicity, cancer (HTP-c) | **HTP-nc**=Human toxicity, non-cancer (HTP-nc) | **IR**=Ionising radiation, human health (IR) | **SQP**=Land use (SQP)

#### CLASSIFICATION OF DISCLAIMERS TO THE DECLARATION OF CORE AND ADDITIONAL ENVIRONMENTAL IMPACT INDICATORS

ILCD classification	Indicator	Disclaimer
ILCD type / level 1	Global warming potential (GWP)	None
	Depletion potential of the stratospheric ozone layer (ODP)	None
	Potential incidence of disease due to PM emissions (PM)	None
	AAcidification potential, Accumulated Exceedance (AP)	None
	Eutrophication potential, Fraction of nutrients reaching freshwater end compartment (EP-freshwater)	None
ILCD type / level 2	Eutrophication potential, Fraction of nutrients reaching marine end compartment (EP-marine)	None
	Eutrophication potential, Accumulated Exceedance (EP-terrestrial)	None
	Formation potential of tropospheric ozone (POCP)	None
	Potential Human exposure efficiency relative to U235 (IRP)	1
	Abiotic depletion potential for non-fossil resources (ADP-minerals&metals)	2
ILCD type / level 3	Abiotic depletion potential for fossil resources (ADP-fossil)	2
	Water (user) deprivation potential, deprivation-weighted water consumption (WDP)	2
	Potential Comparative Toxic Unit for ecosystems (ETP-fw)	2
	Potential Comparative Toxic Unit for humans (HTP-c)	2
	Potential Comparative Toxic Unit for humans (HTP-nc)	2
	Potential Soil quality index (SQP)	2

**Disclaimer 1** – This impact category deals mainly with the eventual impact of low dose ionizing radiation on human health of the nuclear fuel cycle. It does not consider effects due to possible nuclear accidents, occupational exposure nor due to radioactive waste disposal in underground facilities. Potential ionizing radiation from the soil, from radon and from some construction materials is also not measured by this indicator.

**Disclaimer 2** – The results of this environmental impact indicator shall be used with care as the uncertainties on these results are high or as there is limited experienced with the indicator.

## 3 Results

### CORE ENVIRONMENTAL IMPACT INDICATORS EN15804+A1

Abbreviation	Unit	A1	A2	A3	Total
ADPE	Kg Sb	0.00E+0	0.00E+0	2.28E-4	2.28E-4
GWP	Kg CO2 Equiv.	0.00E+0	0.00E+0	4.42E+0	4.42E+0
ODP	Kg CFC-11 Equiv.	0.00E+0	0.00E+0	7.88E-7	7.88E-7
POCP	Kg Ethene Equiv.	0.00E+0	0.00E+0	3.18E-3	3.18E-3
AP	Kg SO2 Equiv.	0.00E+0	0.00E+0	2.71E-2	2.71E-2
EP	Kg PO43- Equiv.	0.00E+0	0.00E+0	5.88E-3	5.88E-3

**ADPE**=Depletion of abiotic resources-elements | **GWP**=Global warming | **ODP**=Ozone layer depletion | **POCP**=Photochemical oxidants creation | **AP**=Acidification of soil and water | **EP**=Eutrophication

### NATIONAL ANNEX NMD

Abbreviation	Unit	A1	A2	A3	Total
ADPF	Kg Sb	0.00E+0	0.00E+0	2.68E-2	2.68E-2
HTP	kg 1.4 DB	0.00E+0	0.00E+0	2.13E+0	2.13E+0
FAETP	kg 1.4 DB	0.00E+0	0.00E+0	3.15E-2	3.15E-2
MAETP	kg 1.4 DB	0.00E+0	0.00E+0	1.05E+2	1.05E+2
TETP	kg 1.4 DB	0.00E+0	0.00E+0	1.32E-2	1.32E-2

**ADPF**=Depletion of abiotic resources-fossil fuels | **HTP**=Human toxicity | **FAETP**=Ecotoxicity. fresh water | **MAETP**=Ecotoxicity. marine water (MAETP) | **TETP**=Ecotoxicity. terrestrie

## 3.2 INDICATORS DESCRIBING RESOURCE USE AND ENVIRONMENTAL INFORMATION BASED ON LIFE CYCLE INVENTORY (LCI)

### PARAMETERS DESCRIBING RESOURCE USE

Abbreviation	Unit	A1	A2	A3	Total
PERE	MJ	0.00E+0	0.00E+0	4.40E+0	4.40E+0

**PERE**=renewable primary energy ex. raw materials | **PERM**=renewable primary energy used as raw materials | **PERT**=renewable primary energy total | **PENRE**=non-renewable primary energy ex. raw materials | **PENRM**=non-renewable primary energy used as raw materials | **PENRT**=non-renewable primary energy total | **SM**=use of secondary material | **RSF**=use of renewable secondary fuels | **NRSF**=use of non-renewable secondary fuels | **FW**=use of net fresh water

### 3 Results

Abbreviation	Unit	A1	A2	A3	Total
PERM	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
PERT	MJ	0.00E+0	0.00E+0	5.47E+0	5.47E+0
PENRE	MJ	0.00E+0	0.00E+0	7.93E+1	7.93E+1
PENRM	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
PENRT	MJ	0.00E+0	0.00E+0	9.13E+1	9.13E+1
SM	Kg	0.00E+0	0.00E+0	0.00E+0	0.00E+0
RSF	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
NRSF	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
FW	M3	0.00E+0	0.00E+0	2.66E-2	2.66E-2

**PERE**=renewable primary energy ex. raw materials | **PERM**=renewable primary energy used as raw materials | **PERT**=renewable primary energy total | **PENRE**=non-renewable primary energy ex. raw materials | **PENRM**=non-renewable primary energy used as raw materials | **PENRT**=non-renewable primary energy total | **SM**=use of secondary material | **RSF**=use of renewable secondary fuels | **NRSF**=use of non-renewable secondary fuels | **FW**=use of net fresh water

#### OTHER ENVIRONMENTAL INFORMATION DESCRIBING WASTE CATEGORIES

Abbreviation	Unit	A1	A2	A3	Total
HWD	Kg	0.00E+0	0.00E+0	1.72E-4	1.72E-4
NHWD	Kg	0.00E+0	0.00E+0	6.02E-1	6.02E-1
RWD	Kg	0.00E+0	0.00E+0	7.01E-4	7.01E-4

**HWD**=hazardous waste disposed | **NHWD**=non hazardous waste disposed | **RWD**=radioactive waste disposed

#### ENVIRONMENTAL INFORMATION DESCRIBING OUTPUT FLOWS

Abbreviation	Unit	A1	A2	A3	Total
CRU	Kg	0.00E+0	0.00E+0	0.00E+0	0.00E+0
MFR	Kg	0.00E+0	0.00E+0	0.00E+0	0.00E+0
MER	Kg	0.00E+0	0.00E+0	0.00E+0	0.00E+0

**CRU**=Components for re-use | **MFR**=Materials for recycling | **MER**=Materials for energy recovery | **EE**=Exported energy | **EET**=Exported Energy Thermic | **EEE**=Exported Energy Electric

### 3 Results

Abbreviation	Unit	A1	A2	A3	Total
EE	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
EET	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0
EEE	MJ	0.00E+0	0.00E+0	0.00E+0	0.00E+0

**CRU**=Components for re-use | **MFR**=Materials for recycling | **MER**=Materials for energy recovery | **EE**=Exported energy | **EET**=Exported Energy Thermic | **EEE**=Exported Energy Electric

## 3 Results

### 3.3 INFORMATION ON BIOGENIC CARBON CONTENT PER TON

#### BIOGENIC CARBON CONTENT

The following Information describes the biogenic carbon content in (the main parts of) the product at the factory gate per ton:

Biogenic carbon content	Amount	Unit
Biogenic carbon content in the product	0	kg C
Biogenic carbon content in accompanying packaging	0	kg C

## 3 Results

### 3.4 ENVIRONMENTAL COST INDICATOR NL PER TON

Using the environmental cost indicator (ECI) method, which is presented in the NMD Determination Method (2020), the results are aggregated to the single-point score. The ECI is a relevant valuation method, especially in the Dutch construction sector. In the Netherlands, it is a prerequisite for public tenders. The aim of the indicator is to show the shadow price for environmental impacts of a product or project. The application of single-point scores is an additional assessment tool for eco-balance results. However, it must be pointed out that weightings are always based on a value maintenance and not on a scientific basis (EN 14040). The ECI results are shown in the following table.

Module EN15804	ECI NL	Share in total (%)
A1 Raw Materials Supply	€ 0,00	0,0 %
A2 Transport	€ 0,00	0,0 %
A3 Manufacturing	€ 0,60	100,0 %
<b>ECI NL per functional unit</b>	<b>€ 0,60</b>	

## 4 Contact information

Publisher	Operator	Owner of declaration
 De Hoop Bouwgrondstoffen B.V. Duitslandweg 2 4538 BK Terneuzen, NL  E-mail: <a href="mailto:info@dehoop-bouwgrondstoffen.nl">info@dehoop-bouwgrondstoffen.nl</a> Website: <a href="http://www.dehoop-bouwgrondstoffen.nl">www.dehoop-bouwgrondstoffen.nl</a>	 De Hoop Bouwgrondstoffen B.V. Duitslandweg 2 4538 BK Terneuzen, NL  E-mail: <a href="mailto:info@dehoop-bouwgrondstoffen.nl">info@dehoop-bouwgrondstoffen.nl</a> Website: <a href="http://www.dehoop-bouwgrondstoffen.nl">www.dehoop-bouwgrondstoffen.nl</a>	 De Hoop Bouwgrondstoffen B.V. Duitslandweg 2 4538 BK Terneuzen, NL  E-mail: <a href="mailto:info@dehoop-bouwgrondstoffen.nl">info@dehoop-bouwgrondstoffen.nl</a> Website: <a href="http://www.dehoop-bouwgrondstoffen.nl">www.dehoop-bouwgrondstoffen.nl</a>